Alternation in the Gap-junctional Intercellular Communication Capacity during the Maturation of Osteocytes in the Embryonic Chick Calvaria




Ziyi Wang, Naoya Odagaki, Tomoyo Tanaka, Mana Hashimoto, Masahiro Nakamura, Satoru Hayano, Yoshihito Ishihara, Noriaki Kawanabe, Hiroshi Kamioka

Department of Orthodontics, Okayama University Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences, Okayama, Japan

Introduction

To further understand the mechanism of bone development, it is therefore important to quantify the difference in Gap-Junctional Intercellular Communication (GJIC) capacity between young and developmentally mature osteocytes. For this aim, we established an embryonic chick calvaria growth model, which reflected the growth of the calvaria in embryos at 13 to 21 days of age. This model can also reveal the relationship between the osteocyte location and its age. So this model actually quantified and confirmed our previous finding that different aged osteocytes have different morphologies, with the young osteocytes having a more spherical shape and developmentally mature osteocytes being more spindle shaped. Next, the FRAP technique was applied to calculate the dye displacement ratio in cell bodies and permeability in the cellular process network. In addition, a comparison was made between the young and developmentally mature osteocytes.



- Conclusions
- 2. GJIC between osteocytes could be regulated and almost half of the gap junction channels have the same permeability as the cell processes for Calcein transfer. Scan it
- Indirect evidence of the active transduction of molecules through the Cx43 channel.

1. GJIC was decreased during the developmental maturation of osteocytes.

Acknowledgements

This work was partially supported by the Japan Society for the Promotion of Science in the form of a Grant-in-Aid for Scientific Research (#16H05549).

for more deta